

DUALITY AND PRODUCTS IN ALGEBRAIC (CO)HOMOLOGY THEORIES

NIELS KOWALZIG AND ULRICH KRÄHMER

ABSTRACT. The origin and interplay of products and dualities in algebraic (co)homology theories is ascribed to a $\times_A$ -Hopf algebra structure on the relevant universal enveloping algebra. This provides a unified treatment for example of results by Van den Bergh about Hochschild (co)homology and by Huebschmann about Lie-Rinehart (co)homology.

1. INTRODUCTION

Most classical (co)homology theories of algebraic objects such as groups or Lie, Lie-Rinehart or associative algebras can be realised as

$$(1) \quad H^\bullet(X, M) := \text{Ext}_U^\bullet(A, M), \quad H_\bullet(X, N) := \text{Tor}_\bullet^U(N, A)$$

for an augmented ring $X = (U, A)$ (a ring with a distinguished left module) that is functorially attached to a given object. The cohomology coefficients are left U -modules M and those in homology are right U -modules N .

Our aim here is to clarify the origin and interplay of multiplicative structures and dualities between such (co)homology groups, and to provide a unified treatment of results by Van den Bergh on Hochschild (co)homology [26] and by Huebschmann on Lie-Rinehart (co)homology [8]. The key concept involved is that of a $\times_A$ -Hopf algebra introduced by Schauenburg [22].

The main results can be summarised as follows:

THEOREM 1. *For any A -biprojective $\times_A$ -Hopf algebra U there is a functor*

$$\otimes : U\text{-}\mathbf{Mod} \times U^{\text{op}}\text{-}\mathbf{Mod} \rightarrow U^{\text{op}}\text{-}\mathbf{Mod}$$

that induces for $M \in U\text{-}\mathbf{Mod}$, $N \in U^{\text{op}}\text{-}\mathbf{Mod}$ and $m, n \geq 0$ natural products

$$\frown : \text{Ext}_U^m(A, M) \times \text{Tor}_n^U(N, A) \rightarrow \text{Tor}_{n-m}^U(M \otimes N, A).$$

If $A \in U\text{-}\mathbf{Mod}$ admits a finitely generated projective resolution of finite length and there exists $d \geq 0$ with $\text{Ext}_U^m(A, U) = 0$ for $m \neq d$, then there is a canonical element

$$[\omega] \in \text{Tor}_d^U(A^*, A), \quad A^* := \text{Ext}_U^d(A, U)$$

such that for $m \geq 0$ and $M \in U\text{-}\mathbf{Mod}$ with $\text{Tor}_q^A(M, A^) = 0$ for $q > 0$*

$$\cdot \frown [\omega] : \text{Ext}_U^m(A, M) \rightarrow \text{Tor}_{d-m}^U(M \otimes A^*, A)$$

is an isomorphism.

As we will recall below, $\times_A$ -bialgebras and $\times_A$ -Hopf algebras generalise bialgebras and Hopf algebras towards possibly noncommutative base algebras A . Besides Hopf algebras, both the universal enveloping algebra

$U(A, L)$ of a Lie-Rinehart algebra (A, L) and the enveloping algebra $A^e = A \otimes_k A^{\text{op}}$ of an associative algebra A are $\times_A$ -Hopf algebras, see Section 2.5.

For any $\times_A$ -bialgebra U , the base algebra A carries a left U -action and the category $U\text{-}\mathbf{Mod}$ of left U -modules is monoidal with unit object A . But only for $\times_A$ -Hopf algebras one has a canonical operation $\otimes$ as in Theorem 1 which turns $U^{\text{op}}\text{-}\mathbf{Mod}$ into a module category over $(U\text{-}\mathbf{Mod}, \otimes, A)$ (Lemma 16).

Any $\times_A$ -Hopf algebra carries two left and two right actions of the base algebra, all commuting with each other. The biprojectivity assumed in Theorem 1 refers to the projectivity of two of these, see Section 2.1. Under this condition, we can use the elegant formalism of suspended monoidal categories from [24] to define for $M, N \in U\text{-}\mathbf{Mod}$ and $P \in U^{\text{op}}\text{-}\mathbf{Mod}$ products

$$\smile : H^m(X, M) \times H^n(X, N) \rightarrow H^{m+n}(X, M \otimes N),$$

$$\frown : H^n(X, N) \times H_p(X, P) \rightarrow H_{p-n}(X, N \otimes P),$$

where we again use the abbreviations from (1) (cf. Sections 3.2 and 3.5).

In the last part of Theorem 1, $A^* = H^d(X, U) = \text{Ext}_U^d(A, U)$ is a right U -module via right multiplication in U , and if we define the functor

$$\hat{} : U\text{-}\mathbf{Mod} \rightarrow U^{\text{op}}\text{-}\mathbf{Mod}, \quad M \mapsto \hat{M} := M \otimes A^*,$$

then the statement can be rewritten as an isomorphism

$$H^m(X, M) \simeq H_{\dim(X)-m}(X, \hat{M}), \quad \dim(X) := \text{proj.dim}_U(A)$$

that is given as in topology by the cap product with the fundamental class $[\omega] \in H_{\dim(X)}(\hat{A})$ which corresponds under the duality to $\text{id}_A \in H^0(A) = \text{Hom}_U(A, A)$. For $M = A$ this simply means that the $H^\bullet(A)$ -module $H_\bullet(A^*)$ is free with generator $[\omega]$.

Theorem 1 is well-known in group and Lie algebra (co)homology. For $U = A \otimes_k A^{\text{op}}$ it reduces to Van den Bergh's result [26] that stimulated a lot of recent research, see e.g. [2, 4, 5, 14]. Note that we do not need Van den Bergh's invertibility assumption about A^* , which says that $\hat{}$ is an equivalence. However, it is satisfied for many well-behaved algebras [ibid.] and implies the condition $\text{Tor}_q^A(M, A^*) = 0$ for arbitrary A -bimodules M (since invertible bimodules are finitely generated projective as one-sided modules from either side). For Lie-Rinehart algebras (A, L) , Theorem 1 is due to Huebschmann [8], and we find the general setting helpful for example to understand the different roles of left and right modules that he has observed. As Huebschmann has showed, the conditions of Theorem 1 are satisfied whenever L is finitely generated projective over A , and A^* coincides as an A -module with $\Lambda_A^d L$ and is in particular projective, so also here we have $\text{Tor}_q^A(M, A^*) = 0$ for arbitrary (A, L) -modules M .

We were also motivated by the current discussion of the numerous bialgebroid generalisations of Hopf algebras, see [1]. Several authors have raised the question where Lie-Rinehart algebras fit in. They were shown in [28, 18] to be $\times_A$ -bialgebras, see also [12, 9]. Here we remark that they are in fact always $\times_A$ -Hopf algebras, but not necessarily Hopf algebroids in the sense of Böhm and Szlachányi (Example 8, this answers a question of Böhm [1]). So both these examples and the applications in homological algebra clearly demonstrate the relevance of the intermediate concept of a $\times_A$ -Hopf algebra.

Theorem 1 could be generalised to differentially graded $\times_A$ -Hopf algebras, sheaves of such, or suitable abstract monoidal categories. One can also drop the condition $\text{Ext}_U^n(A, U) = 0$ for $n \neq d$ and the assumption that $\text{Tor}_q^A(M, A^*) = 0$. Then one obtains for a bounded below chain complex M over $U\text{-}\mathbf{Mod}$ an isomorphism $\text{RHom}_U(A, M) \simeq (M \otimes_A^L \text{RHom}_U(A, U)) \otimes_U^L A$.

N.K. is supported by the NWO through the GQT cluster. U.K. is supported by the EPSRC fellowship EP/E/043267/1 and partially by the Polish Government Grant N201 1770 33. We thank Andy Baker, Gabriella Böhm, Ken Brown, Henning Krause and Valery Lunts for discussions.

2. PRELIMINARIES ON $\times_A$ -HOPF ALGEBRAS

2.1. Some conventions. Throughout this paper, “ring” means “unital and associative ring”, and we fix a commutative ring k . All other algebras, modules etc. will have an underlying structure of a k -module.

Secondly, we fix a k -algebra A , i.e., a ring with a ring homomorphism $\eta_A : k \rightarrow Z(A)$ to its centre. We denote by $A\text{-}\mathbf{Mod}$ the category of left A -modules, by A^{op} the opposite and by $A^e := A \otimes_k A^{\text{op}}$ the enveloping algebra of A . Thus left A^e -modules are A -bimodules with symmetric action of k .

Our main object is finally an algebra U over A^e , where we now refer to the less standard notion of an algebra over a possibly noncommutative base algebra: U is a k -algebra with a k -algebra homomorphism $\eta = \eta_U : A^e \rightarrow U$. This gives rise to a forgetful functor $U\text{-}\mathbf{Mod} \rightarrow A^e\text{-}\mathbf{Mod}$ using which we consider every U -module M also as an A -bimodule with actions

$$(2) \quad a \triangleright m \triangleleft b := \eta(a \otimes_k b)m, \quad a, b \in A, m \in M.$$

Similarly, every right U -module N is also an A -bimodule via

$$(3) \quad a \blacktriangleright m \blacktriangleleft b := n\eta(b \otimes_k a), \quad a, b \in A, n \in N.$$

In particular, U itself carries two left and two right A -actions all commuting with each other. Usually we consider U as an A^e -module using $a \triangleright u \triangleleft b$, and otherwise we write e.g. $\blacktriangleright U \blacktriangleleft$ to denote which actions are considered. Since this will be repeatedly a necessary technical condition, we define:

DEFINITION 2. *For an A^e -algebra U we call $M \in U\text{-}\mathbf{Mod}$ A -biprojective if both $\triangleright M \in A\text{-}\mathbf{Mod}$ and $M \triangleleft \in A^{\text{op}}\text{-}\mathbf{Mod}$ are projective modules.*

2.2. $\times_A$ -bialgebras [25]. Consider an A^e -algebra U as above which is also a coalgebra in the monoidal category $A^e\text{-}\mathbf{Mod}$. That is, there are maps

$$\Delta : U \rightarrow U \otimes_A U, \quad \varepsilon : U \rightarrow A$$

satisfying the usual coalgebra axioms (see e.g. [1] for the details), where

$$(4) \quad U \otimes_A U = U \otimes_k U / \text{span}_k \{u \triangleleft a \otimes_k v - u \otimes_k a \triangleright v \mid a \in A, u, v \in U\}.$$

For $A = k$ one calls U a bialgebra if Δ and ε are algebra homomorphisms, but in general there is no natural algebra structure on $U \otimes_A U$. The way out of this problem was found by Takeuchi [25] and involves the embedding

$$(5) \quad \iota : U \times_A U \rightarrow U \otimes_A U,$$

where $U \times_A U$ is the centre of the A -bimodule $\blacktriangleright U \triangleleft \otimes_A \triangleright U \blacktriangleleft$:

$$U \times_A U := \left\{ \sum_i u_i \otimes_A v_i \in U \otimes_A U \mid \sum_i a \blacktriangleright u_i \otimes_A v_i = \sum_i u_i \otimes_A v_i \blacktriangleleft a \right\}.$$

The product of U turns this into an algebra over A^e , with

$$\eta_{U \times_A U} : A^e \rightarrow U \times_A U, \quad a \otimes_k b \mapsto \eta(a \otimes_k 1) \otimes_A \eta(1 \otimes_k b).$$

Similarly, A is an algebra over k , but not over A^e in general. To handle this one needs the canonical map

$$(6) \quad \pi : \text{End}_k(A) \rightarrow A, \quad \varphi \mapsto \varphi(1),$$

and the fact that $\text{End}_k(A)$ is an algebra over A^e , with

$$\eta_{\text{End}_k(A)} : A^e \rightarrow \text{End}_k(A), \quad (\eta_{\text{End}_k(A)}(a \otimes b))(c) := acb.$$

Now it makes sense to require Δ and ε to factor through ι and π :

DEFINITION 3. *A (left) $\times_A$ -bialgebra is an algebra U over A^e together with two homomorphisms $\hat{\Delta} : U \rightarrow U \times_A U$ and $\hat{\varepsilon} : U \rightarrow \text{End}_k(A)$ of algebras over A^e such that U is a coalgebra in $A^e\text{-}\mathbf{Mod}$ via $\Delta = \iota \circ \hat{\Delta}$ and $\varepsilon = \pi \circ \hat{\varepsilon}$.*

So one has for example for any $\times_A$ -bialgebra

$$\Delta(a \triangleright u \triangleleft b) = a \triangleright u_{(1)} \otimes_A u_{(2)} \triangleleft b, \quad \Delta(a \blacktriangleright u \blacktriangleleft b) = u_{(1)} \blacktriangleleft b \otimes_A a \blacktriangleright u_{(2)},$$

where we started to use Sweedler's shorthand notation $u_{(1)} \otimes_A u_{(2)}$ for $\Delta(u)$.

Be aware that the four A -actions are not the only feature of $\times_A$ -bialgebras that disappears for $A = k$. Another crucial one is for example that the counit $\varepsilon : U \rightarrow A$ is not necessarily a ring homomorphism. Note also that many authors write $s(a) := \eta(a \otimes 1)$ and $t(a) := \eta(1 \otimes a)$ and formulate the theory using these so-called source and target maps rather than η .

2.3. The monoidal category $U\text{-}\mathbf{Mod}$ [21]. Definition 3 might appear complicated, but it is the correct concept from several points of view. An important one for us is the following result of Schauenburg [21, Theorem 5.1]:

THEOREM 4. *The $\times_A$ -bialgebra structures on an algebra $\eta : A^e \rightarrow U$ over A^e correspond bijectively to monoidal structures on $U\text{-}\mathbf{Mod}$ for which the forgetful functor $U\text{-}\mathbf{Mod} \rightarrow A^e\text{-}\mathbf{Mod}$ induced by η is strictly monoidal.*

Given a $\times_A$ -bialgebra structure on U , the monoidal structure on $U\text{-}\mathbf{Mod}$ is defined as for bialgebras: one takes the tensor product $M \otimes_A N$ of the A -bimodules underlying $M, N \in U\text{-}\mathbf{Mod}$ and defines a left U -action via Δ ,

$$(7) \quad u(m \otimes_A n) := u_{(1)}m \otimes_A u_{(2)}n, \quad u \in U, m \in M, n \in N.$$

DEFINITION 5. *If U is a $\times_A$ -bialgebra and $M, N \in U\text{-}\mathbf{Mod}$ are left U -modules, we denote the left U -module $M \otimes_A N$ with U -action (7) by $M \otimes N$.*

The unit object in $U\text{-}\mathbf{Mod}$ is A on which U acts via

$$\hat{\varepsilon}(u)(a) = \varepsilon(a \blacktriangleright u) = \varepsilon(u \blacktriangleleft a),$$

where the last equality is a consequence of the definition of a $\times_A$ -bialgebra.

There is an analogous notion of right $\times_A$ -bialgebra for which $U^{\text{op}}\text{-}\mathbf{Mod}$ is monoidal. However, for a left $\times_A$ -bialgebra there is in general no canonical monoidal structure on $U^{\text{op}}\text{-}\mathbf{Mod}$ or even only right action of U on A .

2.4. **$\times_A$ -Hopf algebras** [22]. Let U be a $\times_A$ -bialgebra and define

$$(8) \quad \beta : \blacktriangleright U \otimes_{A^{\text{op}}} U_{\triangleleft} \rightarrow U_{\triangleleft} \otimes_A \blacktriangleright U, \quad u \otimes_{A^{\text{op}}} v \mapsto u_{(1)} \otimes_A u_{(2)} v,$$

the so-called Galois map of U , where

$$\blacktriangleright U \otimes_{A^{\text{op}}} U_{\triangleleft} = U \otimes_k U / \text{span}\{a \blacktriangleright u \otimes_k v - u \otimes_k v \triangleleft a \mid u, v \in U, a \in A\}.$$

One could flip the tensor components in order to avoid taking the tensor product over A^{op} , but we found it more convenient to keep β in the form which is standard for bialgebras over fields. For the latter it is easily seen that β is bijective if and only if U is a Hopf algebra with $\beta^{-1}(u \otimes_k v) := u_{(1)} \otimes S(u_{(2)})v$, where S is the antipode of U . This motivates the following definition due to Schauenburg [22]:

DEFINITION 6. *A $\times_A$ -bialgebra U is a $\times_A$ -Hopf algebra if β is a bijection.*

Following Schauenburg, we adopt a Sweedler-type notation

$$(9) \quad u_+ \otimes_{A^{\text{op}}} u_- := \beta^{-1}(u \otimes_A 1)$$

for the so-called translation map

$$\beta^{-1}(\cdot \otimes_A 1) : U \rightarrow \blacktriangleright U \otimes_{A^{\text{op}}} U_{\triangleleft}.$$

Since substantial for the subsequent calculations, we list some properties of β^{-1} as proven in [22, Proposition 3.7]: one has for all $u, v \in U, a \in A$

$$(10) \quad u_{+(1)} \otimes_A u_{+(2)} u_- = u \otimes_A 1 \in U_{\triangleleft} \otimes_A \blacktriangleright U$$

$$(11) \quad u_{(1)+} \otimes_{A^{\text{op}}} u_{(1)-} u_{(2)} = u \otimes_{A^{\text{op}}} 1 \in \blacktriangleright U \otimes_{A^{\text{op}}} U_{\triangleleft}$$

$$(12) \quad u_+ \otimes_{A^{\text{op}}} u_- \in U \times_{A^{\text{op}}} U$$

$$(13) \quad u_+ \otimes_{A^{\text{op}}} u_{-(1)} \otimes_A u_{-(2)} = u_{++} \otimes_{A^{\text{op}}} u_- \otimes_A u_{+-}$$

$$(14) \quad (uv)_+ \otimes_{A^{\text{op}}} (uv)_- = u_+ v_+ \otimes_{A^{\text{op}}} v_- u_-$$

$$(15) \quad \eta(a \otimes b)_+ \otimes_{A^{\text{op}}} \eta(a \otimes b)_- = \eta(a \otimes 1) \otimes_{A^{\text{op}}} \eta(b \otimes 1),$$

where in (12) we abbreviated

$$U \times_{A^{\text{op}}} U := \left\{ \sum_i u_i \otimes_{A^{\text{op}}} v_i \in \blacktriangleright U \otimes_{A^{\text{op}}} U_{\triangleleft} \mid \sum_i u_i \triangleleft a \otimes_{A^{\text{op}}} v_i = \sum_i u_i \otimes_{A^{\text{op}}} a \blacktriangleright v_i \right\}$$

and in (13) the tensor product over A^{op} links the first and third tensor component (cf. [22, Equation (3.7)]). By (10) and (12) one can write

$$(16) \quad \beta^{-1}(u \otimes_A v) = u_+ \otimes_{A^{\text{op}}} u_- v$$

which is easily checked to be well-defined over A with (14) and (15).

2.5. **Examples.** Clearly, Hopf algebras over k such as universal enveloping algebras of Lie algebras or group algebras are $\times_k$ -Hopf algebras. But also the enveloping algebra of an associative algebra that governs Hochschild (co)homology is an example as pointed out by Schauenburg [22]:

EXAMPLE 7. The enveloping algebra $U := A^e$ of any k -algebra A is a $\times_A$ -bialgebra with $\eta = \text{id}_{A^e}$ and coproduct and counit

$$\Delta : U \rightarrow U \otimes U, \quad a \otimes_k b \mapsto (a \otimes_k 1) \otimes_A (1 \otimes_k b), \quad \varepsilon : U \rightarrow A, \quad a \otimes_k b \mapsto ab.$$

As for the $\times_A$ -Hopf algebra structure, the tensor product in question reads

$$\blacktriangleright U \otimes_{A^{\text{op}}} U_{\triangleleft} = U \otimes_k U / \text{span}_k \{(a \otimes_k cb) \otimes_k (a' \otimes_k b') - (a \otimes_k b) \otimes_k (a' \otimes_k b'c)\},$$

where cb and $b'c$ is understood to be the product in A . One then easily verifies that

$$(a \otimes_k b)_+ \otimes_{A^{\text{op}}} (a \otimes_k b)_- := (a \otimes_k 1) \otimes_{A^{\text{op}}} (b \otimes_k 1)$$

yields an inverse of the Galois map defined as in (16).

Finally we discuss Lie-Rinehart algebras which define for example Poisson (co)homology. Several authors [28, 12, 18] have shown that their enveloping algebras are $\times_A$ -bialgebras, but they are in fact also $\times_A$ -Hopf algebras:

EXAMPLE 8. Let (A, L) be a Lie-Rinehart algebra over k [20, 6]. We denote by $(a, X) \mapsto aX$ the A -module structure on L and by $(X, a) \mapsto X(a)$ the L -action on A given by the anchor $\hat{\varepsilon} : L \rightarrow \text{Der}_k(A)$. Its universal enveloping algebra $U = U(A, L)$ is the universal k -algebra equipped with two maps

$$\iota_A : A \rightarrow U, \quad \iota_L : L \rightarrow U$$

of k -algebras and of k -Lie algebras, respectively, and subject to the identities

$$\iota_A(a)\iota_L(X) = \iota_L(aX), \quad \iota_L(X)\iota_A(a) - \iota_A(a)\iota_L(X) = \iota_A(X(a))$$

for $a \in A, X \in L$; confer [20] for the precise construction. The map ι_A is injective, so we refrain from further mentioning it. We will also merely write X when we mean $\iota_L(X)$ (if L is A -projective, then ι_L is injective as well).

Recall now from e.g. [28, 18] that U carries the structure of a $\times_A$ -bialgebra: the maps $\eta(- \otimes 1)$ and $\eta(1 \otimes -)$ are equal and given by ι_A . The prescriptions

$$(17) \quad \Delta(X) = 1 \otimes_A X + X \otimes_A 1, \quad \Delta(a) = a \otimes_A 1$$

which map $X \in L$ and $a \in A$ into $U \times_A U$ can be extended by the universal property to a coproduct $\hat{\Delta} : U \rightarrow U \times_A U$. The counit is similarly given by the extension of the anchor $\hat{\varepsilon}$ to U . The bijectivity of the Galois map is seen in the same way: the translation map is given on generators as

$$(18) \quad a_+ \otimes_{A^{\text{op}}} a_- := a \otimes_{A^{\text{op}}} 1, \quad X_+ \otimes_{A^{\text{op}}} X_- := X \otimes_{A^{\text{op}}} 1 - 1 \otimes_{A^{\text{op}}} X.$$

These maps stay in $U \times_{A^{\text{op}}} U$ which is an algebra through the product of U in the first and its opposite in the second tensor factor. By universality we obtain a map $U \rightarrow U \times_{A^{\text{op}}} U$, and then β^{-1} is defined using (16).

On the other hand, U is not necessarily a Hopf algebroid in the sense of [1] (this also answers Böhm's question asked therein whether any $\times_A$ -Hopf algebra is a Hopf algebroid). This structure assumes the existence of an antipode $S : U \rightarrow U^{\text{op}}$ satisfying certain axioms. As a result, the left U -action on A yields by composition with S also a right U -module structure. However, there might be an obstruction for this. For example, take $L = \Gamma(T^{1,0}S^2)$, where $T^{1,0}S^2 \oplus T^{0,1}S^2 = TS^2 \otimes \mathbb{C}$ is the decomposition of the complexified tangent bundle of the 2-sphere $S^2 \subset \mathbb{R}^3$ into the holomorphic and antiholomorphic part with respect to the standard complex structure. This defines together with $A = C^\infty(S^2, \mathbb{C})$ a Lie-Rinehart algebra, where the action of L on A is the usual action of a vector field on a smooth function and the action of A on L is given by fibrewise multiplication. We know by work of Huebschmann [7] that the right U -module structures on A correspond bijectively to left U -module structures on L itself (in general on its top exterior power over A , but here this is L because $T^{1,0}S^2$ is only a line bundle). Such a left U -action corresponds precisely to a flat connection

∇ on the complex line bundle $T^{1,0}S^2$, with $X \in L$ acting on sections of $T^{1,0}S^2$ by the covariant derivative ∇_X (see [7] for the details). But the curvature of any connection represents the first Chern class of the bundle which is nonvanishing since $T^{1,0}S^2$ is not trivial. Therefore, there is no flat connection aka left U -action on L and hence no right U -action on A .

3. MULTIPLICATIVE STRUCTURES

3.1. $\mathcal{D}^-(U)$ as a suspended monoidal category [24]. For any ring U , we denote by $\mathcal{D}^-(U)$ the derived category of bounded above cochain complexes of left U -modules. As usual, we identify any $M \in U\text{-}\mathbf{Mod}$ with a complex in $\mathcal{D}^-(U)$ concentrated in degree 0, and any bounded below chain complex $P_\bullet$ with a bounded above cochain complex by putting $P^n := P_{-n}$.

If U is an A -biprojective $\times_A$ -bialgebra, then any projective $P \in U\text{-}\mathbf{Mod}$ is A -biprojective. Hence the monoidal structure of $U\text{-}\mathbf{Mod}$ extends to a monoidal structure on $\mathcal{D}^-(U)$ with unit object still given by A and product being the total tensor product $\otimes^L = \otimes_A^L$ (the A -biprojectivity of U -projectives is needed for example to have [27, Lemma 10.6.2]).

Together with the shift functor $T : \mathcal{D}^-(U) \rightarrow \mathcal{D}^-(U)$, $(TC)^n = C^{n+1}$, $\mathcal{D}^-(U)$ becomes what is called a suspended monoidal category in [24]. This just means that for all $C, D \in \mathcal{D}^-(U)$, the canonical isomorphisms

$$TC \otimes^L D \simeq T(C \otimes^L D) \simeq C \otimes^L TD$$

given by the obvious renumbering make the diagrams

$$\begin{array}{ccc} A \otimes^L TC & \longrightarrow & TC \\ \downarrow & \swarrow & \\ T(A \otimes^L C) & & \end{array} \quad \begin{array}{ccc} TC \otimes^L A & \longrightarrow & TC \\ \downarrow & \swarrow & \\ T(C \otimes^L A) & & \end{array}$$

commutative and the diagram

$$\begin{array}{ccc} TC \otimes^L TD & \longrightarrow & T(C \otimes^L TD) \\ \downarrow & & \downarrow \\ T(TC \otimes^L D) & \longrightarrow & T^2(C \otimes^L D) \end{array}$$

anticommutative (commutative up to a sign -1).

3.2. $\smile$ and $\circ$ [24]. As a special case of the constructions from [24], we define for any A -biprojective $\times_A$ -bialgebra U and $L, M, N \in U\text{-}\mathbf{Mod}$ the cup product

$$\smile : \text{Ext}_U^m(A, M) \times \text{Ext}_U^n(A, N) \rightarrow \text{Ext}_U^{m+n}(A, M \otimes N)$$

and the classical Yoneda product

$$\circ : \text{Ext}_U^m(N, M) \times \text{Ext}_U^n(L, N) \rightarrow \text{Ext}_U^{m+n}(L, M).$$

The latter is just the composition of morphisms in $\mathcal{D}^-(U)$ if one identifies

$$\text{Ext}_U^n(L, N) \simeq \text{Hom}_{\mathcal{D}^-(U)}(L, T^n N),$$

and

$$\text{Ext}_U^m(N, M) \simeq \text{Hom}_{\mathcal{D}^-(U)}(N, T^m M) \simeq \text{Hom}_{\mathcal{D}^-(U)}(T^n N, T^{m+n} M).$$

The former is obtained as follows: given

$$\varphi \in \text{Ext}_U^m(A, M) \simeq \text{Hom}_{\mathcal{D}^-(U)}(A, T^m M),$$

$$\psi \in \text{Ext}_U^n(A, N) \simeq \text{Hom}_{\mathcal{D}^-(U)}(A, T^n N),$$

one defines $\varphi \smile \psi$ as the composition

$$\begin{aligned} A &\simeq A \otimes A \\ &\xrightarrow{\varphi \otimes \psi} T^m M \otimes^L T^n N \simeq T^m(M \otimes^L T^n N) \simeq T^{m+n}(M \otimes^L N) \\ &\longrightarrow T^{m+n}(M \otimes N), \end{aligned}$$

where the last map is the augmentation $M \otimes^L N \rightarrow H^0(M \otimes^L N) \simeq \text{Tor}_0^A(M, N) \simeq M \otimes N$, or rather T^{m+n} applied to this morphism in $\mathcal{D}^-(U)$.

A straightforward extension of Theorem 1.7 from [24] now gives:

THEOREM 9. *If U is an A -biprojective $\times_A$ -bialgebra, then we have*

$$\psi \circ \varphi = \varphi \smile \psi = (-1)^{mn} \psi \smile \varphi, \quad \varphi \in \text{Ext}_U^m(A, A), \psi \in \text{Ext}_U^n(A, M)$$

as elements of $\text{Ext}_U^{m+n}(A, M) \simeq \text{Ext}_U^{m+n}(A, A \otimes M) \simeq \text{Ext}_U^{m+n}(A, M \otimes A)$.

In particular, $\text{Ext}_U(A, A)$ becomes through either of the products a graded commutative algebra over the commutative subring $\text{Hom}_U(A, A)$.

Proof. This is proven exactly as in [24]. For the reader's convenience we include one of the diagrams involved. The unlabeled arrows are canonical maps coming from the suspended monoidal structure.

$$\begin{array}{ccccc} A & \xrightarrow{\quad} & A \otimes A & & \\ \downarrow \varphi & & \downarrow \text{id} \otimes \varphi & \searrow \psi \otimes \text{id} & \\ T^m A & \xrightarrow{\quad} & A \otimes T^m A & & T^n M \otimes^L A \\ & \searrow & \downarrow & \searrow \psi \otimes \text{id} & \downarrow \text{id} \otimes \varphi \\ & & T^m(A \otimes A) & & T^n M \otimes^L T^m A \\ \downarrow \text{id} & & \downarrow T^m(\psi \otimes \text{id}) & \swarrow & \downarrow \\ T^m A & & T^m(T^n M \otimes^L A) & & T^n(M \otimes^L T^m A) \\ \downarrow T^m(\psi) & \swarrow & \downarrow & \swarrow & \\ T^{m+n} M & \longleftarrow & T^{m+n}(M \otimes^L A) & & \end{array}$$

The morphism $\psi \circ \varphi \in \text{Hom}_{\mathcal{D}^-(U)}(A, T^{m+n} M)$ is the path going straight down from A to $T^{m+n} M$, and $\psi \smile \varphi$ is the one which goes clockwise round the whole diagram. All faces of the diagram commute except the lower right square which introduces a sign $(-1)^{mn}$, so we get $\psi \circ \varphi = (-1)^{mn} \psi \smile \varphi$. The other identity is shown with a similar diagram. $\square$

3.3. Tensoring projectives. This paragraph is a small excursus about the projectivity of the tensor product of two projective objects of a monoidal category. For example, $U \otimes U \in U\text{-}\mathbf{Mod}$ is not necessarily projective even for a bialgebra U over a field $A = k$ (so the A -projectivity of U or the exactness of $\otimes$ does not help). Here is a simple example (for a detailed study of examples of categories of Mackey functors see [15]):

EXAMPLE 10. Consider the bialgebra $U = \mathbb{C}[a, b, c]$ over $A = k = \mathbb{C}$ with

$$\begin{aligned}\Delta(a) &= a \otimes a, & \Delta(b) &= a \otimes b + b \otimes c, & \Delta(c) &= c \otimes c, \\ \varepsilon(a) &= 1, & \varepsilon(b) &= 0, & \varepsilon(c) &= 1.\end{aligned}$$

Geometrically, this is the coordinate ring of the complex algebraic semigroup G of upper triangular 2×2 -matrices, and Δ and ε are dual to the semigroup law $G \times G \rightarrow G$ and the embedding of the identity matrix into G .

We prove that $U \otimes U \in U\text{-}\mathbf{Mod}$ is not projective by considering the fibres of the semigroup law $G \times G \rightarrow G$. The fibre over a generic and hence invertible element is 3-dimensional, but over 0 it is 4-dimensional, and this will imply our claim. We can use for example [16, Theorem 19 on p. 79]:

THEOREM 11. *Let $U \subset V$ be a flat extension of commutative Noetherian rings, $\mathfrak{p} \subset V$ be a prime ideal and $\mathfrak{q} := U \cap \mathfrak{p}$. Then*

$$\dim(V_{\mathfrak{p}}) = \dim(U_{\mathfrak{q}}) + \dim(V_{\mathfrak{p}} \otimes_U U(\mathfrak{q})),$$

where $\dim$ denotes the Krull dimension of a ring, $V_{\mathfrak{p}}$ is the localisation of V at $\mathfrak{p}$ and $U(\mathfrak{q}) := U_{\mathfrak{q}}/\mathfrak{q}U_{\mathfrak{q}}$ is the residue field of the localisation $U_{\mathfrak{q}}$.

Apply this to our example $U \simeq \Delta(U) \subset V := U \otimes U$: let $\mathfrak{p}$ be the ideal of V generated by $a \otimes_{\mathbb{C}} 1, 1 \otimes_{\mathbb{C}} a, b \otimes_{\mathbb{C}} 1, 1 \otimes_{\mathbb{C}} b, c \otimes_{\mathbb{C}} 1, 1 \otimes_{\mathbb{C}} c$. Geometrically, V is the coordinate ring of $\mathbb{C}^6$ and $V_{\mathfrak{p}}$ is the local ring in 0, so $\dim(V_{\mathfrak{p}}) = 6$. Since $1 \notin \mathfrak{p}$, $\mathfrak{q} = U \cap \mathfrak{p}$ is proper, and it contains the ideal generated by $\Delta(a) = a \otimes_{\mathbb{C}} a, \Delta(b) = a \otimes_{\mathbb{C}} b + b \otimes_{\mathbb{C}} c, \Delta(c) = c \otimes_{\mathbb{C}} c$ which is maximal in U , so $\mathfrak{q} \subset U$ is the ideal generated by a, b, c , and $U_{\mathfrak{q}}$ is the local ring of $\mathbb{C}^3$ at 0 with $\dim(U_{\mathfrak{q}}) = 3$. The field $U(\mathfrak{q})$ is obviously $\mathbb{C}$, and we can write $V_{\mathfrak{p}} \otimes_U U(\mathfrak{q})$ also as $V_{\mathfrak{p}}/\Delta(\mathfrak{q})V_{\mathfrak{p}}$. Since $\Delta(\mathfrak{q})V_{\mathfrak{p}}$ is contained in the ideal $\mathfrak{r}$ generated in $V_{\mathfrak{p}}$ by the elements $a \otimes_{\mathbb{C}} 1, 1 \otimes_{\mathbb{C}} c$, we have $\dim(V_{\mathfrak{p}}/\Delta(\mathfrak{q})V_{\mathfrak{p}}) \geq \dim(V_{\mathfrak{p}}/\mathfrak{r})$. Now $V_{\mathfrak{p}}/\mathfrak{r}$ is the local ring of $\mathbb{C}^4 \subset \mathbb{C}^6$ at 0 and hence $\dim(V_{\mathfrak{p}}/\mathfrak{r}) = 4$. In total, we get the strict inequality $3 + \dim(V_{\mathfrak{p}}/\Delta(\mathfrak{q})V_{\mathfrak{p}}) \geq 3 + 4 = 7 > 6$, and hence V is not flat over U and in particular not projective.

For $\times_A$ -Hopf algebras the situation is, however, much simpler: notice that

$$\blacktriangleright U \otimes_{A^{\text{op}}} M \triangleleft := U \otimes_k M / \text{span}\{a \blacktriangleright u \otimes_k m - u \otimes_k m \triangleleft a \mid u \in U, a \in A, m \in M\}$$

is for any $\times_A$ -bialgebra U and $M \in U\text{-}\mathbf{Mod}$ a left U -module by left multiplication on the first factor. Just as for $M = U$, there is a Galois map

$$\beta_M : \blacktriangleright U \otimes_{A^{\text{op}}} M \triangleleft \rightarrow U \otimes M, \quad u \otimes_{A^{\text{op}}} m \mapsto u_{(1)} \otimes_A u_{(2)} m,$$

and we have:

LEMMA 12. *For any $\times_A$ -bialgebra U , the generalised Galois map β_M is a morphism of U -modules. If U is a $\times_A$ -Hopf algebra, then β_M is bijective.*

Proof. The U -linearity of β_M follows immediately from the fact that $\hat{\Delta} : U \rightarrow U \times_A U$ is a homomorphism of algebras over A^e . Furthermore, if β is a bijection, then β_M is so as well since we can identify β_M with $\beta \otimes_U \text{id}_M$, and then the inverse is simply given by $\beta_M^{-1}(u \otimes_A m) = u_+ \otimes_{A^{\text{op}}} u_- m$. $\square$

Using this one now gets:

THEOREM 13. *If U is a $\times_A$ -Hopf algebra and $U_{\triangleleft} \in A^{\text{op}}\text{-Mod}$ is projective, then $P \otimes Q \in U\text{-Mod}$ is projective for all projectives $P, Q \in U\text{-Mod}$.*

Proof. By assumption, any projective module over U is also projective over A^{op} , and if $\varphi : R \rightarrow S$ is any ring map, then $S \otimes_R : R\text{-Mod} \rightarrow S\text{-Mod}$ maps projectives to projectives. This shows that $\blacktriangleright U \otimes_{A^{\text{op}}} U_{\triangleleft}$ and hence (Lemma 12) $U \otimes U$ is projective. Since $\otimes = \otimes_A$ commutes with arbitrary direct sums, $P \otimes Q$ is projective for all projectives P, Q . $\square$

COROLLARY 14. *If U is as in Theorem 13 and $P \in \mathcal{D}^-(U)$ is a projective resolution of $A \in U\text{-Mod}$, then so is $P \otimes P := \text{Tot}(P_{\bullet} \otimes P_{\bullet}) = P \otimes^L P$.*

This leads to the traditional construction of $\smile$ given for $A = k$ in [3, Chapter XI]: one fixes a projective resolution P of A , and by the above, $\text{Ext}_U(A, M \otimes N)$ is the total (co)homology of the double (cochain) complex

$$C_{mn}^2 := \text{Hom}_U(P_m \otimes P_n, M \otimes N).$$

Then $\smile$ is given as the composition of the canonical map

$$\begin{aligned} & \bigoplus_{m+n=p} \text{Ext}_U^m(A, M) \otimes_k \text{Ext}_U^n(A, N) \\ & \simeq \bigoplus_{m+n=p} H^m(\text{Hom}_A(P_{\bullet}, M)) \otimes_k H^n(\text{Hom}_A(P_{\bullet}, N)) \\ & \rightarrow H^p\left(\bigoplus_{m+n=\bullet} \text{Hom}_A(P_m, M) \otimes_k \text{Hom}_A(P_n, N)\right) = H^p(\text{Tot}(C_{\bullet\bullet}^1)) \end{aligned}$$

where $C_{mn}^1 := \text{Hom}_U(P_m, M) \otimes_k \text{Hom}_U(P_n, N)$, with the map

$$H(\text{Tot}(C_{\bullet\bullet}^1)) \rightarrow H(\text{Tot}(C_{\bullet\bullet}^2)) \simeq \text{Ext}_U(A, M \otimes N)$$

that is induced by the morphism of double complexes

$$C_{mn}^1 \ni \varphi \otimes_k \psi \mapsto \{x \otimes y \mapsto \varphi(x) \otimes \psi(y)\} \in C_{mn}^2.$$

For the sake of completeness let us finally remark that as for $A = k$ one can in particular use the bar construction to obtain a canonical resolution:

LEMMA 15. *For any $\times_A$ -bialgebra U , the complex of left U -modules*

$$C_n^{\text{bar}} := (\blacktriangleright U_{\triangleleft})^{\otimes_{A^{\text{op}}} n+1}, \quad u(v_0 \otimes_{A^{\text{op}}} \cdots \otimes_{A^{\text{op}}} v_n) := uv_0 \otimes_{A^{\text{op}}} \cdots \otimes_{A^{\text{op}}} v_n$$

whose boundary map is given by

$$\begin{aligned} b' : u_0 \otimes_{A^{\text{op}}} \cdots \otimes_{A^{\text{op}}} u_n & \mapsto \sum_{i=0}^{n-1} (-1)^i u_0 \otimes_{A^{\text{op}}} \cdots \otimes_{A^{\text{op}}} u_i u_{i+1} \otimes_{A^{\text{op}}} \cdots \otimes_{A^{\text{op}}} u_n \\ & \quad + (-1)^n u_0 \otimes_{A^{\text{op}}} \cdots \otimes_{A^{\text{op}}} \varepsilon(u_n) \blacktriangleright u_{n-1} \end{aligned}$$

is a contractible resolution of $A \in U\text{-Mod}$ with augmentation

$$\varepsilon : C_0^{\text{bar}} = U \rightarrow A =: C_{-1}^{\text{bar}},$$

and if $U_{\triangleleft} \in A^{\text{op}}\text{-}\mathbf{Mod}$ is projective, then $C_n^{\text{bar}} \in U\text{-}\mathbf{Mod}$ is projective.

Proof. All claims are straightforward: there is a contracting homotopy

$$s : C_n^{\text{bar}} \rightarrow C_{n+1}^{\text{bar}}, \quad u_0 \otimes_{A^{\text{op}}} \cdots \otimes_{A^{\text{op}}} u_n \mapsto 1 \otimes_{A^{\text{op}}} u_0 \otimes_{A^{\text{op}}} \cdots \otimes_{A^{\text{op}}} u_n, \quad n \geq 0,$$

$$s : A = C_{-1}^{\text{bar}} \rightarrow U = C_0^{\text{bar}}, \quad a \mapsto \eta(a \otimes 1),$$

and the projectivity of C_n^{bar} follows as in the proof of Theorem 13. $\square$

3.4. The functor $\otimes : U\text{-}\mathbf{Mod} \times U^{\text{op}}\text{-}\mathbf{Mod} \rightarrow U^{\text{op}}\text{-}\mathbf{Mod}$. Now we introduce the functor $\otimes$ mentioned in Theorem 1.

LEMMA 16. *Let U be a $\times_A$ -Hopf algebra and $M \in U\text{-}\mathbf{Mod}$, $P \in U^{\text{op}}\text{-}\mathbf{Mod}$ be left and right U -modules, respectively. Then the formula*

$$(19) \quad (m \otimes_A p)u := u_- m \otimes_A p u_+, \quad u \in U, \quad m \in M, p \in P,$$

defines a right U -module structure on the tensor product

$$(20) \quad M \otimes_A P := M \otimes_k P / \text{span}\{m \triangleleft a \otimes_k p - m \otimes_k a \blacktriangleright p \mid a \in A\}.$$

If N is any other (left) U -module, then the canonical isomorphism

$$(21) \quad (M \otimes N) \otimes_A P \simeq M \otimes_A (N \otimes_A P)$$

of A -bimodules is also an isomorphism in $U^{\text{op}}\text{-}\mathbf{Mod}$. Finally, the tensor flip

$$(M \otimes_A P) \otimes_U N \rightarrow P \otimes_U (N \otimes_A M), \quad m \otimes_A p \otimes_U n \mapsto p \otimes_U n \otimes_A m$$

is an isomorphism of k -modules.

Proof. To show firstly that (19) is well-defined over A , we compute

$$\begin{aligned} (m \otimes_A (a \blacktriangleright p))u &= u_- m \otimes_A p \eta(1 \otimes a)u_+ = u_- m \otimes_A p(u_+ \triangleleft a) \\ &= (a \blacktriangleright u_-)m \otimes_A p u_+ = u_- (\eta(1 \otimes a)m) \otimes_A p u_+ \\ &= ((m \triangleleft a) \otimes_A p)u, \end{aligned}$$

where (12) and the action properties were used. Together with (20) this also proves the well-definedness with respect to the presentation of $u_+ \otimes_{A^{\text{op}}} u_-$. With the help of (14) one sees immediately that for $u, v \in U$ we have

$$(m \otimes_A p)(uv) = (uv)_- m \otimes_A p(uv)_+ = v_- u_- m \otimes_A p u_+ v_+ = ((m \otimes_A p)u)v,$$

since P and M were right and left U -modules, respectively. As a conclusion, $M \otimes_A P \in U^{\text{op}}\text{-}\mathbf{Mod}$. Equation (21) is a direct consequence of the associativity of the tensor product of A -bimodules and of (13).

For the last part one has to check that the flip is well-defined: we have

$$\begin{aligned} \eta(1 \otimes a)m \otimes_A p \otimes_U n &\mapsto p \otimes_U n \otimes_A \eta(1 \otimes a)m = p \otimes_U \eta(1 \otimes a)(n \otimes_A m) \\ &= p \eta(1 \otimes a) \otimes_U (n \otimes_A m), \end{aligned}$$

which is what $m \otimes_A p \eta(1 \otimes a) \otimes_U n$ gets mapped to. And secondly, we have

$$\begin{aligned} m \otimes_A p \otimes_U un &\mapsto p \otimes_U un \otimes_A m = p \otimes_U (u_+)_{(1)} n \otimes_A (u_+)_{(2)} u_- m \\ &= p \otimes_U u_+(n \otimes_A u_- m) = p u_+ \otimes_U n \otimes_A u_- m, \end{aligned}$$

which is what $u_- m \otimes_A p u_+ \otimes_U n = (m \otimes_A p)u \otimes_U n$ gets mapped to. $\square$

DEFINITION 17. *We denote the above constructed U^{op} -module by $M \otimes P$.*

Thus an unadorned $\otimes$ refers from now on either to the monoidal product on $U\text{-}\mathbf{Mod}$ or to the just defined action of $U\text{-}\mathbf{Mod}$ on $U^{\text{op}}\text{-}\mathbf{Mod}$. For example, (21) would now simply be written as $(M \otimes N) \otimes P \simeq M \otimes (N \otimes P)$.

EXAMPLE 18. Let (A, L) be a Lie-Rinehart algebra and M be a left and N a right $U(A, L)$ -module, respectively (or, in the terminology of [6, 8], left and right (A, L) -modules). Using (18), one gets the right $U(A, L)$ -module structure on $M \otimes_A N$ from formula (2.4) in [8, p. 112]:

$$(m \otimes_A n)X = m \otimes_A nX - Xm \otimes_A n, \quad m \in M, n \in N, X \in L.$$

If we assume again that U is A -biprojective, then the above results extend directly to the derived category $\mathcal{D}^-(U^{\text{op}})$: we obtain a functor

$$\otimes^L = \otimes_A^L : \mathcal{D}^-(U) \times \mathcal{D}^-(U^{\text{op}}) \rightarrow \mathcal{D}^-(U^{\text{op}})$$

and we have for all $M, N \in \mathcal{D}^-(U)$, $P \in \mathcal{D}^-(U^{\text{op}})$ canonical isomorphisms

$$(22) \quad (M \otimes^L N) \otimes^L P \simeq M \otimes^L (N \otimes^L P), \quad (M \otimes^L P) \otimes_U^L N \simeq P \otimes_U^L (N \otimes^L M).$$

3.5. $\frown$ and $\bullet$. These products are dual to $\smile$ and $\circ$. The first one is

$$\bullet : \text{Ext}_U^m(L, M) \times \text{Tor}_n^U(N, L) \rightarrow \text{Tor}_{n-m}^U(N, M)$$

which exists for a ring U and $L, M \in U\text{-}\mathbf{Mod}$, $N \in U^{\text{op}}\text{-}\mathbf{Mod}$: an element

$$\varphi \in \text{Ext}_U^m(L, M) \simeq \text{Hom}_{\mathcal{D}^-(U)}(L, T^m M)$$

defines a morphism in $\mathcal{D}^-(\mathbb{Z})$

$$N \otimes_U^L L \rightarrow N \otimes_U^L T^m M, \quad x \otimes_U y \mapsto x \otimes_U \varphi(y),$$

and $\varphi \bullet \cdot$ is the induced map in (co)homology

$$\begin{aligned} \text{Tor}_n^U(N, L) &\simeq H^{-n}(N \otimes_U^L L) \\ &\xrightarrow{H^{-n}(\text{id} \otimes \varphi)} H^{-n}(N \otimes_U^L T^m M) \simeq H^{m-n}(N \otimes_U^L M) \simeq \text{Tor}_{n-m}^U(N, M). \end{aligned}$$

For $M \in U\text{-}\mathbf{Mod}$, $N \in U^{\text{op}}\text{-}\mathbf{Mod}$ as before, the cap product

$$\frown : \text{Ext}_U^m(A, M) \times \text{Tor}_n^U(N, A) \rightarrow \text{Tor}_{n-m}^U(M \otimes N, A)$$

involves the functor $\otimes$ from the previous paragraph, so for this we want U to be an A -biprojective $\times_A$ -Hopf algebra again. Similarly as for $\bullet$,

$$\varphi \in \text{Ext}_U^m(A, M) \simeq \text{Hom}_{\mathcal{D}^-(U)}(A, T^m M)$$

defines a morphism in $\mathcal{D}^-(k)$

$$\begin{aligned} N \otimes_U^L A &\simeq N \otimes_U^L (A \otimes A) \\ &\xrightarrow{\text{id} \otimes \text{id} \otimes \varphi} N \otimes_U^L (A \otimes^L T^m M) \simeq N \otimes_U^L (T^m A \otimes^L M) \simeq (M \otimes^L N) \otimes_U^L T^m A \\ &\longrightarrow (M \otimes N) \otimes_U^L T^m A, \end{aligned}$$

where the last $\simeq$ in the second line is induced by the tensor flip as in the derived version (22) of Lemma 16, and the morphism from the second to the third line is similarly as in the definition of $\smile$ induced by the morphism $M \otimes^L N \rightarrow M \otimes N$ in $\mathcal{D}^-(U^{\text{op}})$ that takes zeroth cohomology. Passing now to cohomology we get $\varphi \frown \cdot : \text{Tor}_n^U(N, A) \rightarrow \text{Tor}_{n-m}^U(M \otimes N, A)$.

More explicitly, if $P \in \mathcal{D}^-(U)$ is a projective resolution of A , then $\cap$ is induced by the morphism

$$B_{ij}^1 \ni n \otimes_U (x \otimes_A y) \mapsto \{\varphi \mapsto (\varphi(y) \otimes_A n) \otimes_U x\} \in B_{ij}^2$$

from the double complex

$$B_{ij}^1 := N \otimes_U (P_i \otimes_A P_j)$$

whose total homology is $\mathrm{Tor}^U(N, A)$ to the double complex

$$B_{ij}^2 := \mathrm{Hom}_k(\mathrm{Hom}_U(P_j, M), (M \otimes N) \otimes_U P_i)$$

whose homology has a natural map to $\mathrm{Hom}_k(\mathrm{Ext}_U(A, M), \mathrm{Tor}^U(M \otimes N, A))$.

In direct analogy with Theorem 9 we get:

THEOREM 19. *If U is an A -biprojective $\times_A$ -Hopf algebra, then we have*

$$\varphi \bullet (x \otimes_U y) = \varphi \cap (x \otimes_U y), \quad \varphi \in \mathrm{Ext}_U^m(A, A), x \otimes_U y \in N \otimes_U^L A$$

as elements of $N \otimes_U^L A \simeq (A \otimes N) \otimes_U^L A$.

4. DUALITY AND THE PROOF OF THEOREM 1

4.1. The underived case. In the special case that A is finitely generated projective itself, Theorem 1 reduces to standard linear algebra. We go through this case first since it is both instructive and used in the proof of the general case. For the reader's convenience we include full proofs.

LEMMA 20. *Let U be a ring, $A \in U\text{-}\mathbf{Mod}$ be finitely generated projective, and A^* be $\mathrm{Hom}_U(A, U)$ with its canonical U^{op} -module structure.*

1. *A^* is finitely generated projective, and if $e_1, \dots, e_n$ are generators of A , then there exist generators $e^1, \dots, e^n \in A^*$ with*

$$\sum_i e^i(a) e_i = a, \quad \sum_i e^i \alpha(e_i) = \alpha$$

for all $a \in A$ and $\alpha \in A^*$. The element

$$\omega := \sum_i e^i \otimes e_i \in A^* \otimes_U A$$

is independent of the choice of the generators e_i, e^j .

2. *For all U^{op} -modules M , the assignment*

$$\delta(m \otimes a)(\alpha) := m\alpha(a), \quad m \in M, a \in A, \alpha \in A^*$$

extends uniquely to an isomorphism of abelian groups

$$\delta : M \otimes_U A \rightarrow \mathrm{Hom}_{U^{\mathrm{op}}}(A^*, M).$$

3. *One has $(A^*)^* \simeq A$ and $A^* \otimes_U M \simeq \mathrm{Hom}_U(A, M)$ for $M \in U\text{-}\mathbf{Mod}$.*

Proof. Since A is projective, there is a splitting $\iota : A \rightarrow U^n$ of

$$\pi : U^n \rightarrow A, \quad (u_1, \dots, u_n) \mapsto \sum_i u_i e_i.$$

Hence $U^n \simeq A \oplus A_\perp$ for some $A_\perp \in U\text{-}\mathbf{Mod}$. Dually this gives $A^* \oplus (A_\perp)^* = (U^n)^* \simeq U^n$, whence A^* is finitely generated projective. The e^i

can be defined as the composition of ι with the projection of U^n on the i -th summand. This proves the first parts of 1. For 2. just note that

$$\mathrm{Hom}_{U^{\mathrm{op}}}(A^*, M) \ni \varphi \mapsto \sum_i \varphi(e^i) \otimes e_i \in M \otimes_U A$$

inverts δ . Since $\omega = \delta^{-1}(\mathrm{id}_{A^*})$, it does indeed not depend on the choice of generators. 3. now follows from 1. and 2. $\square$

As in the introduction, let us abbreviate in the situation of this theorem

$$H^0(M) := \mathrm{Hom}_U(A, M), \quad H_0(N) := N \otimes_U A$$

for $M \in U\text{-}\mathbf{Mod}$, $N \in U^{\mathrm{op}}\text{-}\mathbf{Mod}$, and call $\omega \in H_0(A^*)$ the fundamental class of (U, A) . Then 3. says for $M = A$ that we have an isomorphism

$$(23) \quad \cdot \bullet \omega : H^0(A) \rightarrow H_0(A^*), \quad \varphi \mapsto \sum_i e^i \otimes \varphi(e_i).$$

Using Lemma 16 we can upgrade this to the underived case of Theorem 1:

LEMMA 21. *Let U be a $\times_A$ -Hopf algebra and assume A is finitely generated projective as a U -module. Then the cap product with the fundamental class $\omega \in H_0(A^*) = A^* \otimes_U A$ defines for all $M \in U\text{-}\mathbf{Mod}$ an isomorphism*

$$\cdot \frown \omega : H^0(M) \rightarrow H_0(M \otimes A^*).$$

Proof. We have $\varphi \frown \omega = \sum_i (\varphi(1) \otimes_A e^i) \otimes_U e_i$, and Lemma 16 identifies

$$H_0(M \otimes A^*) = (M \otimes A^*) \otimes_U A \simeq A^* \otimes_U (A \otimes M) \simeq A^* \otimes_U M.$$

In this chain of identifications, $\varphi \frown \omega$ is mapped to

$$\varphi \frown \omega \mapsto \sum_i e^i \otimes_U (e_i \otimes_A \varphi(1)) \mapsto \sum_i e^i \otimes_U (e_i \varphi(1)) = \sum_i e^i \otimes_U \varphi(e_i)$$

which is identified with φ under the isomorphism $\mathrm{Hom}_U(A, M) \simeq A^* \otimes_U M$ given by $\varphi \mapsto \sum_i e^i \otimes_U \varphi(e_i)$ as in (23). The claim follows. $\square$

4.2. The derived case. It remains to throw in some homological algebra to obtain Theorem 1 in general. To shorten the presentation, we define:

DEFINITION 22. *A module A over a ring U is perfect if it admits a finite resolution by finitely generated projectives. We call such a module a duality module if there exists $d \geq 0$ such that $\mathrm{Ext}_U^n(A, U) = 0$ for all $n \neq d$. We abbreviate in this case $A^* := \mathrm{Ext}_U^d(A, U)$ and call d the dimension of A .*

The main remaining step is to prove a derived version of Lemma 20. One could use a result of Neeman by which $A \in U\text{-}\mathbf{Mod}$ is perfect if and only if $\mathrm{Hom}_U(A, \cdot)$ commutes with direct sums [11, 19], or the Ischebeck spectral sequence which degenerates at E^2 if A is a duality module [10, 13, 23]. However, we include a more elementary and self-contained proof.

THEOREM 23. *Let $A \in U\text{-}\mathbf{Mod}$ be a duality module of dimension d .*

1. *The projective dimension of $A \in U\text{-}\mathbf{Mod}$ is d .*
2. *A^* is a duality module of the same dimension d .*

3. If $P_\bullet \rightarrow A$ is a finitely generated projective resolution of length d , then $P_{d-\bullet}^* = \text{Hom}_U(P_{d-\bullet}, U)$ is a finitely generated projective resolution of A^* and the canonical isomorphism

$$\delta : M \otimes_U P_i \rightarrow \text{Hom}_U(P_i^*, M), \quad m \otimes_U p \mapsto \{\alpha \mapsto m\alpha(p)\}$$

induces for all U^{op} -modules M a canonical isomorphism

$$\text{Tor}_i^U(M, A) \rightarrow \text{Ext}_{U^{\text{op}}}^{d-i}(A^*, M).$$

4. There is a canonical isomorphism $(A^*)^* \simeq A$.

Proof. Let $P_\bullet \rightarrow A$ be a finitely generated projective resolution of finite length $m \geq 0$ (which exists since A is perfect). Then the (co)homology of

$$0 \rightarrow P_0^* \rightarrow \dots \rightarrow P_m^* \rightarrow 0, \quad P_n^* = \text{Hom}_U(P_n, U)$$

is $\text{Ext}_U^\bullet(A, U)$, so by assumption we have $m \geq d$ and the above complex is exact except at P_d^* where the homology is $A^* = \text{Ext}_U^d(A, U)$. Furthermore, all the P_n^* are finitely generated projective since the P_n are so (Lemma 20).

Let π_i be the map $P_i^* \rightarrow P_{i+1}^*$ and put $K := \ker \pi_{d+1}$. By construction,

$$(24) \quad 0 \rightarrow K \rightarrow P_{d+1}^* \rightarrow \dots \rightarrow P_m^* \rightarrow 0$$

is exact. If one compares this exact sequence with the sequence

$$\dots \rightarrow 0 \rightarrow 0 \rightarrow P_m^* \rightarrow P_m^* \rightarrow 0$$

using Schanuel's lemma (see [17, 7.1.2]), one obtains that K is projective.

The exactness of $P_\bullet^*$ at P_{d+1}^* gives $K = \text{im } \pi_d$, and by the projectivity of K , the map $\pi_d : P_d^* \rightarrow K \subset P_{d+1}^*$ splits so that $P_d^* \simeq K \oplus K_\perp$, $K_\perp := \ker \pi_d$. In particular, both K and $K_\perp$ are finitely generated.

It follows from all this that the complex

$$(25) \quad 0 \rightarrow P_0^* \rightarrow \dots \rightarrow P_{d-1}^* \rightarrow K_\perp \rightarrow 0$$

is a finitely generated projective resolution of A^* : since $\text{im } \pi_{d-1} \subset P_d^*$ is contained in $\ker \pi_d = K_\perp$, the above complex is still exact at P_{d-1}^* , and the homology at $K_\perp$ is the homology of $P_\bullet^*$ at P_d^* , that is, A^* .

Since (24) is a finitely generated projective resolution of 0 and $P_{d-\bullet}^*$ is as a complex a direct sum of (25) and (a shift of) (24) we also know that $\text{Ext}_{U^{\text{op}}}^\bullet(A^*, M)$ is for any $M \in U^{\text{op}}\text{-Mod}$ the (co)homology of $\text{Hom}_U(P_{d-\bullet}^*, M)$. By Lemma 20, this is isomorphic as a chain complex to $M \otimes_U P_{d-\bullet}$ via the isomorphism given in 3., and the homology of this complex is $\text{Tor}_{d-\bullet}^U(M, A)$. This proves 3. The special case $M = U$ implies the remaining claims. $\square$

Assume finally that in the situation of the above theorem, U is an A -biprojective $\times_A$ -Hopf algebra. Since P is a projective resolution, we have $M \otimes_U P \simeq M \otimes_U^L P$ and $\text{Hom}_U(P^*, M) \simeq \text{RHom}_U(P^*, M)$, and δ gives an isomorphism between them. The fundamental class is defined to be

$$\omega := \delta^{-1}(\text{id}_{A^*}) \in A^* \otimes_U^L A \simeq P^* \otimes_U A \simeq A^* \otimes_U P,$$

and Theorem 23 gives immediately:

COROLLARY 24. *If $e_1, \dots, e_n$ and $\tilde{e}^1, \dots, \tilde{e}^n$ are generators of A and of A^* , respectively, then there are $e^1, \dots, e^n \in P_0^*$ and $\tilde{e}_1, \dots, \tilde{e}_n \in P_d$ such that*

$$\omega = \sum_i e^i \otimes_U e_i = \sum_i \tilde{e}^i \otimes_U \tilde{e}_i,$$

and δ is given by the Yoneda product $\cdot \bullet \omega$.

Theorem 1 follows now as in the underived case (Lemma 21) working with $\mathrm{RHom}_U(A, M)$ and $(M \otimes^L A^*) \otimes_U^L A$ instead of $H^0(M) = \mathrm{Hom}_U(A, M)$ and $H_0(M \otimes A^*) = (M \otimes A^*) \otimes_U A$: using Theorem 19 and (22) one gets

$$\begin{aligned} (M \otimes^L A^*) \otimes_U^L A &\simeq A^* \otimes_U^L (A \otimes^L M) \simeq A^* \otimes_U^L M \\ &\simeq P^* \otimes_U^L M \simeq \mathrm{RHom}_U(P, M) \\ &\simeq \mathrm{RHom}_U(A, M), \end{aligned}$$

where we hide the reindexing of the complexes for the sake of better readability (so P^* stands for $P_{d-\bullet}^*$, and $\mathrm{RHom}_U(P, M)$ and $\mathrm{RHom}_U(A, M)$ are reindexed in the same way). This leads to a convergent spectral sequence

$$\mathrm{Tor}_p^U(\mathrm{Tor}_q^A(M, A^*), A) \Rightarrow \mathrm{Ext}_U^{d-p-q}(A, M),$$

and under the last assumption of Theorem 1 ($\mathrm{Tor}_q^A(M, A^*) = 0$ for $q > 0$) this spectral sequence degenerates to the claimed isomorphism.

REFERENCES

- [1] Gabriella Böhm, *Hopf algebroids*, (2008), preprint [arXiv:0805.3806v1](#).
- [2] Kenneth A. Brown and James J. Zhang, *Dualising complexes and twisted Hochschild (co)homology for Noetherian Hopf algebras*, J. Algebra **320** (2008), no. 5, 1814–1850.
- [3] Henri Cartan and Samuel Eilenberg, *Homological algebra*, Princeton University Press, Princeton, N. J., 1956.
- [4] Vasilij Dolgushev, *The Van den Bergh duality and the modular symmetry of a Poisson variety*, (2006), preprint [arXiv:math/0612288](#).
- [5] Victor Ginzburg, *Calabi-Yau algebras*, (2006), preprint [arXiv:math/0612139](#).
- [6] Johannes Huebschmann, *Poisson cohomology and quantization*, J. Reine Angew. Math. **408** (1990), 57–113.
- [7] ———, *Lie-Rinehart algebras, Gerstenhaber algebras and Batalin-Vilkovisky algebras*, Ann. Inst. Fourier (Grenoble) **48** (1998), no. 2, 425–440.
- [8] ———, *Duality for Lie-Rinehart algebras and the modular class*, J. Reine Angew. Math. **510** (1999), 103–159.
- [9] ———, *The universal Hopf algebra associated with a Hopf-Lie-Rinehart algebra*, (2008), preprint [arXiv:0802.3836](#).
- [10] Friedrich Ischebeck, *Eine Dualität zwischen den Funktoren Ext und Tor*, J. Algebra **11** (1969), 510–531.
- [11] Bernhard Keller, *On differential graded categories*, International Congress of Mathematicians. Vol. II, Eur. Math. Soc., Zürich, 2006, pp. 151–190.
- [12] Masoud Khalkhali and Bahram Rangipour, *Cyclic cohomology of (extended) Hopf algebras*, Noncommutative geometry and quantum groups (Warsaw, 2001), Banach Center Publ., vol. 61, Polish Acad. Sci., Warsaw, 2003, pp. 59–89.
- [13] Ulrich Krämer, *Poincaré duality in Hochschild (co)homology*, New techniques in Hopf algebras and graded ring theory, K. Vlaam. Acad. Belgie Wet. Kunsten (KVAB), Brussels, 2007, pp. 117–125.
- [14] Stéphane Launois and Lionel Richard, *Twisted Poincaré duality for some quadratic Poisson algebras*, Lett. Math. Phys. **79** (2007), no. 2, 161–174.
- [15] L. Gaunce Lewis, Jr., *When projective does not imply flat, and other homological anomalies*, Theory Appl. Categ. **5** (1999), No. 9, 202–250 (electronic).

- [16] Hideyuki Matsumura, *Commutative algebra*, second ed., Mathematics Lecture Note Series, vol. 56, Benjamin/Cummings Publishing Co., Inc., Reading, Mass., 1980.
- [17] John C. McConnell and James C. Robson, *Noncommutative Noetherian rings*, revised ed., Graduate Studies in Mathematics, vol. 30, American Mathematical Society, Providence, RI, 2001, With the cooperation of L. W. Small.
- [18] Ieke Moerdijk and Janez Mrčun, *On the universal enveloping algebra of a Lie-Rinehart algebra*, (2008), preprint [arXiv:0801.3929](https://arxiv.org/abs/0801.3929).
- [19] Amnon Neeman, *Triangulated categories*, Annals of Mathematics Studies, vol. 148, Princeton University Press, Princeton, NJ, 2001.
- [20] George S. Rinehart, *Differential forms on general commutative algebras*, Trans. Amer. Math. Soc. **108** (1963), 195–222.
- [21] Peter Schauenburg, *Bialgebras over noncommutative rings and a structure theorem for Hopf bimodules*, Appl. Categ. Structures **6** (1998), no. 2, 193–222.
- [22] ———, *Duals and doubles of quantum groupoids ($\times_R$ -Hopf algebras)*, New trends in Hopf algebra theory (La Falda, 1999), Contemp. Math., vol. 267, Amer. Math. Soc., Providence, RI, 2000, pp. 273–299.
- [23] Evgenij G. Sklyarenko, *Poincaré duality and relations between the functors Ext and Tor* , Mat. Zametki **28** (1980), no. 5, 769–776, 803.
- [24] Mariano Suarez-Alvarez, *The Hilton-Heckmann argument for the anti-commutativity of cup products*, Proc. Amer. Math. Soc. **132** (2004), no. 8, 2241–2246 (electronic).
- [25] Mitsuhiro Takeuchi, *Groups of algebras over $A \otimes \overline{A}$* , J. Math. Soc. Japan **29** (1977), no. 3, 459–492.
- [26] Michel Van den Bergh, *A relation between Hochschild homology and cohomology for Gorenstein rings*, Proc. Amer. Math. Soc. **126** (1998), no. 5, 1345–1348, Erratum: Proc. Amer. Math. Soc. **130**, no. 9, 2809–2810 (electronic) (2002).
- [27] Charles A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994.
- [28] Ping Xu, *Quantum groupoids*, Comm. Math. Phys. **216** (2001), no. 3, 539–581.

N.K.: UTRECHT UNIVERSITY, DEPARTMENT OF MATHEMATICS, P.O. BOX 80.010,
3508TA UTRECHT, THE NETHERLANDS
E-mail address: nkowalzi@science.uva.nl

U.K.: UNIVERSITY OF GLASGOW, DEPARTMENT OF MATHEMATICS, UNIVERSITY GAR-
DENS, GLASGOW G12 8QW, SCOTLAND
E-mail address: ukraehmer@maths.gla.ac.uk